

---

**MATH 135 Fall 2020: Extra Practice 10**
**Warm-Up Exercises**

**WE01.** Express  $\frac{2-i}{3+4i}$  in standard form.

*Solution.* Multiply numerator and denominator by the conjugate of the denominator:

$$\frac{2-i}{3+4i} = \frac{(2-i)(3-4i)}{9+16} = \frac{2-11i}{25} = \frac{2}{25} - \frac{11}{25}i \quad \square$$

**WE02.** Write  $x = \frac{9+i}{5-4i}$  in polar form,  $r(\cos \theta + i \sin \theta)$ , with  $0 \leq \theta < 2\pi$ .

*Solution.* We express first in standard form by multiplying through the conjugate:

$$\frac{9+i}{5-4i} = \frac{(9+i)(5+4i)}{41} = \frac{41+41i}{41} = 1+i$$

We can geometrically interpret this as  $\sqrt{2} \operatorname{cis} \frac{\pi}{4}$ . □

**WE03.** Write  $(\sqrt{3} + i)^4$  in standard form.

*Solution.* We first place the quantity within the brackets in polar form. By inspection, this is  $2 \operatorname{cis} \frac{\pi}{6}$ . Now, applying DMT, we have  $(2 \operatorname{cis} \frac{\pi}{6})^4 = 2^4 \operatorname{cis}^4 \frac{\pi}{6} = 16 \operatorname{cis} \frac{2\pi}{3}$ .

Expressing in standard form,  $16(\cos \frac{2\pi}{3} + i \sin \frac{2\pi}{3}) = 16(-\frac{1}{2} + i\frac{\sqrt{3}}{2}) = -8 + 8\sqrt{3}i$  □

**WE04.** Find all  $z \in \mathbb{C}$  such that  $z^5 = 1$  and plot the solutions in the complex plane. (You may state values in polar form.)

*Solution.* Note that  $1 = 1 \operatorname{cis} 0$ . Applying the CRNT, we have that the five roots are given by  $\sqrt[5]{1} \operatorname{cis} (\frac{2k\pi}{5})$  for  $k = 0, 1, 2, 3, 4$ . These values are  $\{1, \operatorname{cis} \frac{\pi}{5}, \operatorname{cis} \frac{4\pi}{5}, \operatorname{cis} \frac{6\pi}{5}, \operatorname{cis} \frac{8\pi}{5}\}$ . I am too lazy to learn `tikz` to draw the diagram. □

**WE05.** Find all  $z \in \mathbb{C}$  such that  $z^2 = \frac{1+i}{1-i}$ .

*Solution.* Simplifying the fraction on the right-hand side,  $\frac{(1+i)(1+i)}{2} = \frac{1+2i-1}{2} = i$ . On the complex plane,  $i = 1 \operatorname{cis} \frac{\pi}{2}$ . Then, by CRNT, the solutions are  $\operatorname{cis} \frac{\pi}{4}$  and  $\operatorname{cis} \frac{5\pi}{4}$ . Evaluating to get standard form, we have  $z = \pm(\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i)$ . □

**Recommended Problems**

**RP01.** Express the following complex numbers in standard form.

(a)  $\frac{(\sqrt{2} - i)^2}{(\sqrt{2} + i)(1 - \sqrt{2}i)}$

*Solution.* Multiply through conjugates of the denominator:

$$\begin{aligned} \frac{(\sqrt{2}-i)^2}{(\sqrt{2}+i)(1-\sqrt{2}i)} &= \frac{(1-2\sqrt{2}i)(\sqrt{2}-i)(1+\sqrt{2}i)}{(3)(3)} \\ &= -\frac{(5-\sqrt{2}i)(\sqrt{2}-i)}{9} \\ &= -\frac{4\sqrt{2}-7i}{9} \\ &= -\frac{4\sqrt{2}}{9} + \frac{7}{9}i \end{aligned} \quad \square$$

(b)  $(\sqrt{5}-i\sqrt{3})^4$

*Solution.* Let  $z = \sqrt{5} - i\sqrt{3}$ . We have  $z^2 = 5 - 2\sqrt{15}i - 3 = 2 - 2\sqrt{15}i$ . Finally,  $z^4 = (z^2)^2 = 4 - 8\sqrt{15}i - 60 = -56 - 8\sqrt{15}i$ .  $\square$

**RP02.** Prove all of the Properties of Complex Arithmetic that were not proved in the notes or in class.

*Proof.* Let  $u = a + bi$ ,  $v = c + di$ , and  $z = f + gi$  be complex numbers. We must show the Properties of Complex Arithmetic, i.e., that

1. Complex addition is associative.

First,  $u + v = (a + c) + (b + d)i$  and  $(u + v) + z = ((a + c) + f) + ((b + d) + g)i$ . Then,  $v + z = (c + f) + (d + g)i$ , so  $u + (v + z) = (a + (c + f)) + (b + (d + g))i$ . The result follows by the associativity of real addition.

2. Complex addition is commutative.

We have  $u + v = (a + c) + (b + d)i = (c + a) + (d + b)i = v + u$  by the commutativity of real addition.

3. The complex additive identity is  $0 = 0 + 0i$ . (Example 3, p. 159)

4. A complex additive inverse  $-z$  exists. (Example 3, p. 159)

5. Complex multiplication is associative.

By definition,  $uv = (ac - bd) + (ad + bc)i$ , so we have

$$(uv)w = ((ac - bd)f - (ad + bc)g) + ((ac - bd)g + (ad + bc)f)i$$

We also have  $vw = (cf - dg) + (cg + df)i$  and by extension

$$\begin{aligned} u(vw) &= (a(cf - dg) - b(cg + df)) + (a(cg + df) + b(cf - dg))i \\ &= (acf - adg - bcb - bdf) + (acg + adf + bcf - bdg)i \\ &= (acf - bdf - adg - bcb) + (acg - bdg + adf + bcf)i \\ &= ((ac - bd)f - (ad + bc)g) + ((ac - bd)g + (ad + bc)f)i \\ &= (uv)w \end{aligned}$$

as desired.

6. Complex multiplication is commutative.

Again,  $uv = (ac - bd) + (ad + bc)i$  and  $vu = (ca - db) + (cb + da)i$ . The result follows from the commutativity of real multiplication and addition.

7. The complex multiplicative identity is  $1 = 1 + 0i$ . (Example 3, p. 159)

8. A complex multiplicative inverse  $z^{-1}$  exists iff  $z \neq 0$ . (Proposition 1, p. 159)

9. Complex multiplication distributes over addition.

We have  $u + v = (a + c) + (b + d)i$ . Then,

$$z(u + v) = (f(a + c) - g(b + d)) + (f(b + d) + g(a + c))i$$

Now,  $zu = (fa - gb) + (fb + ga)i$  and  $zv = (fc - gd) + (fd + gc)i$ , so by definition,

$$\begin{aligned} zu + zv &= ((fa - gb) + (fb + ga)) + ((fc - gd) + (fd + gc))i \\ &= (fa + fc - gb - gd) + (fb + fd + ga + gc)i \\ &= (f(a + c) - g(b + d)) + (f(b + d) + g(a + c))i \\ &= z(u + v) \end{aligned}$$

completing the proof. □

**RP03.** Let  $n \in \mathbb{N}$ . Prove that if  $n \equiv 1 \pmod{4}$ , then  $i^n = i$ .

*Proof.* Let  $n$  be a natural number congruent to 1 modulo 4. Then, we may write  $n = 4k + 1$  for some integer  $k$ . Notice that  $i^4 = (i^2)^2 = (-1)^2 = 1$ .

Therefore,  $i^{4k+1} = (i^4)^k i^1 = (1)^k i = i$ , as desired. □

**RP04.** Find all  $z \in \mathbb{C}$  which satisfy

(a)  $z^2 + 2z + 1 = 0$

*Solution.* Factor:  $z^2 + 2z + 1 = (z + 1)^2$  so  $z = -1 + 0i$  (by [RP06](#)) □

(b)  $z^2 + 2\bar{z} + 1 = 0$

*Solution.* Let  $z = a + bi$  so  $\bar{z} = a - bi$  for two real numbers  $a$  and  $b$ . Then,

$$\begin{aligned} 0 &= z^2 + 2\bar{z} + 1 \\ 0 &= (a + bi)^2 + 2(a - bi) + 1 \\ 0 &= (a^2 + 2a - b^2 + 1) + (2ab - 2b)i \end{aligned}$$

which is true if and only if both  $a^2 + 2a - b^2 + 1 = 0$  and  $2ab - 2b = 0$ .

The second equation implies  $2ab = 2b$  so  $a = 1$  or  $b = 0$ .

If  $a = 1$  then  $a^2 + 2a - b^2 + 1 = 4 - b^2 = 0$ , so  $b = \pm 2$ .

If  $b = 0$ , then  $a^2 + 2a + 1 = (a + 1)^2 = 0$ , so  $a = -1$ .

Therefore, the solutions are  $-1 + 0i$ ,  $1 + 2i$ , and  $1 - 2i$ . □

(c)  $z^2 = \frac{1+i}{1-i}$

*Solution.* Simplify:  $z^2 = \frac{(1+i)^2}{2} = \frac{2i}{2} = i$ . The square roots of  $i$  are  $\pm(\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i)$ .  $\square$

**RP05.**

(a) Find all  $w \in \mathbb{C}$  satisfying  $w^2 = -15 + 8i$ .

*Solution.* We rewrite  $w = a + bi$  for some reals  $a$  and  $b$ . Then,  $(a + bi)^2 = (a^2 - b^2) + (2ab)i = -15 + 8i$ . Equating real and complex parts,  $a^2 - b^2 = -15$  and  $2ab = 8$ .

Now,  $|w^2| = |ww| = |w||w| = |w|^2$  by PM4. Then,  $a^2 + b^2 = \sqrt{(-15)^2 + (8)^2} = 17$ . Solving the system in  $a^2$  and  $b^2$ ,  $a^2 = 1$  and  $b^2 = 16$ .

Therefore,  $a = \pm 1$  and  $b = \pm 4$ . To satisfy  $2ab = 8$ , we must have  $z = \pm(1 + 4i)$ .  $\square$

(b) Find all  $z \in \mathbb{C}$  satisfying  $z^2 - (3 + 2i)z + 5 + i = 0$ .

*Solution.* We apply the quadratic formula. The discriminant is a solution to  $w^2 = (3 + 2i)^2 - 4(1)(5 + i) = (5 + 12i) - (20 + 4i) = -15 + 8i$ . From above, a solution is  $w = 1 + 4i$ . Therefore, the solutions are  $z = \frac{(3+2i) \pm (1+4i)}{2(1)}$ .

The first is  $z = \frac{(3+2i)+(1+4i)}{2} = 2 + 3i$  and the second is  $z = \frac{(3+2i)-(1+4i)}{2} = 1 - i$ .  $\square$

**RP06.** Let  $z, w \in \mathbb{C}$ . Prove that if  $zw = 0$  then  $z = 0$  or  $w = 0$ .

*Proof.* Let  $z$  and  $w$  be complex numbers such that  $zw = 0$ . Suppose for a contradiction that both  $z$  and  $w$  are non-zero. Then, by PM1,  $|z| \neq 0$  and  $|w| \neq 0$ . However, by PM4,  $|zw| = |z||w| \neq 0$ , which is a contradiction, since  $zw = 0$ .

Therefore,  $z$  or  $w$  is zero.  $\square$

**RP07.** Let  $a, b, c \in \mathbb{C}$ . Prove: if  $|a| = |b| = |c| = 1$ , then  $\overline{a + b + c} = \frac{1}{a} + \frac{1}{b} + \frac{1}{c}$ .

*Proof.* First, consider some arbitrary complex number  $z = a + bi$  with modulus 1. By definition,  $a^2 + b^2 = 1^2 = 1$ . Then,  $z^{-1} = \frac{1}{a+bi} = \frac{a-bi}{(a+bi)(a-bi)} = \frac{a-bi}{1} = a - bi = \bar{z}$

Let  $a, b$ , and  $c$  be complex numbers with modulus 1. From above,  $a^{-1} = \bar{a}$ ,  $b^{-1} = \bar{b}$ , and  $c^{-1} = \bar{c}$ . The conclusion immediately follows from PCJ2:


$$\begin{aligned} \overline{a + b + c} &= \bar{a} + \bar{b} + \bar{c} \\ &= \frac{1}{a} + \frac{1}{b} + \frac{1}{c} \end{aligned} \quad \square$$

**RP08.** Find all  $z \in \mathbb{C}$  satisfying  $z^2 = |z|^2$ .

*Proof.* Let  $z$  be a complex number. Recall that  $|z|^2 = \bar{z}z$  by PM3. Then, we have  $z^2 = \bar{z}z$  so  $z = \bar{z}$ , that is,  $z - \bar{z} = 0$ . By PCJ3, this is true if  $2 \operatorname{Im}(z)i = 0$ , which means that  $z$  is purely real. Therefore,  $z$  is any purely real number.  $\square$

**RP09.** Find all  $z \in \mathbb{C}$  satisfying  $|z + 1|^2 \leq 3$  and shade the corresponding region in the complex plane.

*Solution.* We write  $z = a+bi$ , so  $|z+1|^2 = |(a+1)+bi|^2 = (\sqrt{(a+1)^2 + b^2})^2 = (a+1)^2 + b^2$ . Then, we are shading the inside of the circle defined by  $(a+1)^2 + b^2 = 3$ .



This is the circle centered at  $(-1, 0)$  with radius  $\sqrt{3}$ . □

**RP10.** Let  $z, w \in \mathbb{C}$  such that  $\bar{z}w \neq 1$ . Prove that if  $|z| = 1$  or  $|w| = 1$ , then  $\left| \frac{z-w}{1-\bar{z}w} \right| = 1$ .

*Proof* (by sooshi). Let  $z$  and  $w$  be complex numbers such that  $\bar{z}w \neq 1$ . Suppose that  $|z| = 1$  or  $|w| = 1$ . If  $z = w$  and  $|z| = |w| = 1$ , then  $\bar{z}w = \bar{z}z = |z|^2 = 1$ . Therefore,  $z \neq w$ .

Now, consider the case when  $|z| = 1$ . Then,

$$\left| \frac{z-w}{1-\bar{z}w} \right| = \frac{|z-w|}{|1-\bar{z}w|} = \frac{|z||z-w|}{|z||1-\bar{z}w|} = \frac{(1)|z-w|}{|z-z\bar{z}w|} = \frac{|z-w|}{|z-w|} = 1$$

Likewise, if  $|w| = 1$ , then

$$\left| \frac{z-w}{1-\bar{z}w} \right| = \frac{|z-w|}{|1-\bar{z}w|} = \frac{|z-w|}{|w\bar{w}-\bar{z}w|} = \frac{|z-w|}{|w||\bar{w}-\bar{z}|} = \frac{|z-w|}{|w-z|} = 1$$

since  $|w-z| = |-(z-w)| = |-1||z-w| = |z-w|$ , completing the proof. □

**RP11.** Show that for all complex numbers  $z$ ,  $|\operatorname{Re}(z)| + |\operatorname{Im}(z)| \leq \sqrt{2}|z|$ .

*Proof.* Let  $z = r \operatorname{cis} \theta$  be a complex number. Then,  $|z| = r$ ,  $\operatorname{Re}(z) = r \cos \theta$  and  $\operatorname{Im}(z) = r \sin \theta$ . Due to the symmetry of sine and cosine, instead of taking absolute values, we

restrict without loss of generality to the first quadrant  $0 \leq \theta \leq \frac{\pi}{2}$ . Now,

$$\begin{aligned} \operatorname{Re}(z) + \operatorname{Im}(z) &= r(\cos \theta + \sin \theta) \\ &= r\sqrt{2} \frac{\sqrt{2}}{2} (\cos \theta + \sin \theta) \\ &= r\sqrt{2} \left( \frac{\sqrt{2}}{2} \cos \theta + \frac{\sqrt{2}}{2} \sin \theta \right) \\ &= r\sqrt{2} \left( \sin \frac{\pi}{4} \cos \theta + \cos \frac{\pi}{4} \sin \theta \right) \\ &= r\sqrt{2} \sin \left( \frac{\pi}{4} + \theta \right) \\ &\leq r\sqrt{2}(1) \\ &= \sqrt{2}|z| \end{aligned}$$

completing the proof. □

**RP12.** Use *De Moivre's Theorem* (DMT) to prove that  $\sin 4\theta = 4 \sin \theta \cos^3 \theta - 4 \sin^3 \theta \cos \theta$  for all  $\theta \in \mathbb{R}$ .

*Proof.* Let  $\theta \in \mathbb{R}$  and note that by DMT, we have

$$(\cos \theta + i \sin \theta)^4 = \cos 4\theta + i \sin 4\theta$$

so we may say that  $\sin 4\theta = \operatorname{Im}((\cos \theta + i \sin \theta)^4)$ . Expanding this quantity by hand,

$$\begin{aligned} (\cos \theta + i \sin \theta)^4 &= (\cos^2 \theta + 2i \cos \theta \sin \theta - \sin^2 \theta)^2 \\ &= \cos^4 \theta + \sin^4 \theta - 2 \cos^2 \theta \sin^2 \theta + 4i \cos^3 \theta \sin \theta - 4i \sin^3 \theta \cos \theta \\ &= (\cos^4 \theta - 2 \cos^2 \theta \sin^2 \theta + \sin^4 \theta) + (4 \cos^3 \theta \sin \theta - 4 \sin^3 \theta \cos \theta)i \end{aligned}$$

and we have that

$$\sin 4\theta = \operatorname{Im}((\cos \theta + i \sin \theta)^4) = 4 \cos^3 \theta \sin \theta - 4 \sin^3 \theta \cos \theta$$

as desired. □

**RP13.** Let  $n \in \mathbb{N}$  and  $a, b \in \mathbb{R}$ . Show that  $z = (a + bi)^n + (a - bi)^n$  is real.

*Proof.* Let  $n$  be a natural number and  $u = a + bi$  be a complex number. Then,  $\bar{u} = a - bi$ . It inductively follows from PCJ4 and the associativity of multiplication that  $(\bar{u})^n = \overline{u^n}$ .

Now, the fact that  $z = u^n + \overline{u^n}$  is real follows immediately from PCJ3. □

**RP14.** An  $n$ -th root of unity is any complex solution to  $z^n = 1$ . Prove that if  $w$  is an  $n$ -th root of unity,  $\frac{1}{w}$  is also an  $n$ -th root of unity.

*Proof.* Let  $n$  be a natural number and  $w$  be an  $n$ -th root of unity, so  $w^n = 1$ . Knowing that  $1 = \operatorname{cis} 0$ , the CNRT states that  $w = \operatorname{cis}(\frac{2k\pi}{n})$  for some  $0 \leq k < n$ .

By PMC, notice that  $w \operatorname{cis}(-\frac{2k\pi}{n}) = \operatorname{cis}(\frac{2k\pi}{n} - \frac{2k\pi}{n}) = \operatorname{cis} 0 = 1$ , so  $\operatorname{cis}(-\frac{2k\pi}{n})$  is the multiplicative inverse  $w^{-1}$  of  $w$ . Now, since  $\operatorname{cis}$  is  $2\pi$ -periodic, we have

$$\operatorname{cis} \left( -\frac{2k\pi}{n} \right) = \operatorname{cis} \left( 2\pi - \frac{2k\pi}{n} \right) = \operatorname{cis} \left( \frac{2n\pi - 2k\pi}{n} \right) = \operatorname{cis} \left( \frac{2(n-k)\pi}{n} \right)$$

but since  $0 \leq k < n$ , we also have that  $0 \leq n - k < n$ . Therefore, by the CNRT,  $w^{-1}$  is an  $n$ -th root of unity.  $\square$

**RP15.** A complex number  $z$  is called a *primitive*  $n$ -th root of unity if  $z^n = 1$  and  $z^k \neq 1$  for all  $1 \leq k \leq n - 1$ .

(a) For each  $n = 1, 3, 5, 6$  list all the primitive  $n$ -th roots of unity.

*Solution.* Recall that  $1^x = 1$  for any real  $x$ . Applying the CNRT, there are  $n$   $n$ -th roots of unity, of the form

$$z = \text{cis} \left( \frac{2\pi k}{n} \right)$$

for some integer  $0 \leq k < n$ . Note that 1 is always an  $n$ -th root of unity but only a primitive first root of unity. Therefore, we can ignore the case  $k = 0$ .

The only primitive 1st root of unity is 1.

The primitive 3rd roots of unity are  $\text{cis} \frac{2\pi}{3} = \frac{\sqrt{3}}{2} - \frac{1}{2}i$  and  $\text{cis} \frac{4\pi}{3} = \frac{\sqrt{3}}{2} + \frac{1}{2}i$ .

For this, we remain in polar form as calculating sines and cosines of fractions over 5 is *pain*. The primitive 5th roots of unity are  $\text{cis} 0 = 1$ ,  $\text{cis} \frac{2\pi}{5}$ ,  $\text{cis} \frac{4\pi}{5}$ ,  $\text{cis} \frac{6\pi}{5}$ , and  $\text{cis} \frac{8\pi}{5}$ .

The 6th roots of unity are  $\text{cis} \frac{2\pi k}{6} = \text{cis} \frac{\pi k}{3}$ . However, when  $k = 2$ ,  $k = 3$ , and  $k = 4$ , these are also 2nd/3rd roots of unity. Thus, the primitive roots of unity are  $\text{cis} \frac{\pi}{3} = \frac{1}{2} + \frac{\sqrt{3}}{2}i$  and  $\text{cis} \frac{5\pi}{3} = \frac{1}{2} - \frac{\sqrt{3}}{2}i$ .  $\square$

(b) Let  $z$  be a primitive  $n$ -th root of unity. Prove the following statements:

i. For any  $j \in \mathbb{Z}$ ,  $z^j = 1$  if and only if  $n \mid j$ .

*Proof.* Let  $n$  be a natural number,  $j$  be an integer, and  $z$  be a primitive  $n$ -th root of unity so  $z^n = 1$ . Proceed by mutual implication.

( $\Rightarrow$ ) Suppose  $z^j = 1$ . By the Division Algorithm,  $j = qn + r$  for integers  $q$  and  $0 \leq r < n$ . Then,  $1 = z^j = z^{qn+r} = z^{qn}z^r = (z^n)^qz^r = 1^qz^r = z^r$ .

If  $r = 0$ , then  $j = qn$  and  $n \mid j$ . Otherwise, we have  $1 \leq r \leq n - 1$  and  $z^r = 1$ , which is a contradiction to the fact that  $z$  is a primitive  $n$ -th root of unity.

Therefore,  $r = 0$  and  $n \mid j$ .

( $\Leftarrow$ ) If  $n \mid j$  and  $j = nk$  for an integer  $k$ , then  $z^j = z^{nk} = (z^n)^k = 1^k = 1$ .  $\square$

ii. For any  $m \in \mathbb{Z}$ , if  $\text{gcd}(m, n) = 1$ , then  $z^m$  is a primitive  $n$ -th root of unity.

*Proof* (new and improved by sooshi). Let  $z$  be a primitive  $n$ -th root of unity and  $m$  an integer coprime to  $n$ .

Suppose for a contradiction that  $z^m$  is a  $k$ -th root of unity for some  $1 \leq k < n$ . Then,  $(z^m)^k = z^{mk} = 1$ . From above, this implies that  $n \mid mk$  and by CAD,  $n \mid k$ . However, BBD gives that  $n \leq k$ , which is a contradiction.

Therefore,  $z^m$  is a primitive  $n$ -th root of unity.  $\square$

**RP16.** Let  $u$  and  $v$  be fixed complex numbers. Let  $\omega$  be a non-real cube root of unity. For each  $k \in \mathbb{Z}$ , define  $y_k \in \mathbb{C}$  by the formula

$$y_k = \omega^k u + \omega^{-k} v$$

(a) Compute  $y_1$ ,  $y_2$ , and  $y_3$  in terms of  $u$ ,  $v$ , and  $\omega$ .

*Solution.* From RP15(a), the only real cube root of unity is 1, so  $\omega \neq 1$ . In fact,  $\omega = \text{cis } \frac{n\pi}{3}$  for either  $n = 2$  or  $n = 4$ .

If  $n = 2$ , then  $\omega^{-1} = \text{cis } \frac{-2\pi}{3} = \text{cis } \frac{4\pi}{3}$ . If  $n = 4$ , then  $\omega^{-1} = \text{cis } \frac{-4\pi}{3} = \text{cis } \frac{2\pi}{3}$ .

However, using the standard form from RP15(a),  $\text{cis } \frac{2\pi}{3} = \overline{\text{cis } \frac{4\pi}{3}}$ . Therefore,  $\omega^{-1} = \bar{\omega}$ .

Now,  $y_1 = \omega u + \bar{\omega} v$ ,  $y_2 = \omega^2 u + \bar{\omega}^2 v$ , and  $y_3 = \omega^3 u + \bar{\omega}^3 v = u + v$ .  $\square$

(b) Show that  $y_k = y_{k+3}$  for any  $k \in \mathbb{Z}$ .

*Proof.* Let  $k$  be an integer. Then, knowing that both  $\omega$  and  $\bar{\omega}$  are cube roots of unity,

$$\begin{aligned} y_{k+3} &= \omega^{k+3} u + \bar{\omega}^{k+3} v \\ &= \omega^k \omega^3 u + \bar{\omega}^k \bar{\omega}^3 v \\ &= \omega^k u + \bar{\omega}^k v \\ &= y_k \end{aligned}$$

completing the proof.  $\square$

(c) Show that for any  $k \in \mathbb{Z}$ ,

$$y_k - y_{k+1} = \omega^k (1 - \omega)(u - \omega^{k-1} v)$$

*Proof.* Let  $k$  be an integer. Expand the right-hand side:

$$\begin{aligned} \omega^k (1 - \omega)(u - \omega^{k-1} v) &= (\omega^k - \omega^{k+1})(u - \omega^{k-1} v) \\ &= \omega^k u - \omega^{2k+1} v - \omega^{k+1} u + \omega^{2k+2} v \\ &= (\omega^k u + \omega^{2k+2} v) - (\omega^{k+1} u + \omega^{2k+1} v) \end{aligned}$$

To simplify, we show that  $\omega^{2k+2} = \omega^{-k}$ . Equivalently,  $\omega^{2k+2} \omega^k = \omega^{3k+2} = 1$ . Let  $j = k + 1$ . Then,

$$\omega^{3k+2} = \omega^{3(j-1)+2} = \omega^{3j-1} = (\omega^3)^j \omega^{-1} = 1^j \omega^{-1} = \omega^{-1}$$

as desired. Now, we have  $\omega^{2k+2} = \omega^{-k}$  and  $\omega^{2k+1} = \omega^{-(k+1)}$  so

$$\begin{aligned} \omega^k (1 - \omega)(u - \omega^{k-1} v) &= (\omega^k u + \omega^{2k+2} v) - (\omega^{k+1} u + \omega^{2k+1} v) \\ &= (\omega^k u + \omega^{-k} v) - (\omega^{k+1} u + \omega^{-(k+1)} v) \\ &= y_k - y_{k+1} \end{aligned} \quad \square$$

## Challenges

**C01.** Let  $z, w \in \mathbb{C}$ .

(a) Prove that  $|z + w| \leq |z| + |w|$ .



*Proof.* This is the Triangle Inequality, for which a geometric proof is provided in Chapter 10.3. In short, for complex numbers  $z = a + bi$  and  $w = c + di$ , we consider a triangle  $\triangle OZW$  with points  $O(0, 0)$ ,  $Z(a, b)$ , and  $W(c, d)$  in the complex plane. Then,  $|z| = \ell_{OZ}$ ,  $|w| = \ell_{OW}$ , and  $|z + w| = \ell_{ZW}$ . The length of one side of a triangle cannot exceed the sum of the lengths of the other two sides.

Equivalently,  $\ell_{ZW} \leq \ell_{OZ} + \ell_{OW}$ .  $\square$

(b) Prove that  $||z| - |w|| \leq |z - w| \leq |z| + |w|$ .

*Proof.* Let  $z$  and  $w$  be complex numbers. We prove the inequalities separately.

We apply the Triangle Inequality with  $z$  and  $-w$ . Then,  $|z + (-w)| \leq |z| + |-w|$  but  $|-w| = |-1||w| = |w|$  by PM4, so we have  $|z - w| \leq |z| + |w|$ .

Now, notice that  $|z| = |(z - w) + w| \leq |z - w| + |w|$  so  $|z| - |w| \leq |z - w|$ .

Likewise,  $|w| = |(w - z) + z| \leq |w - z| + |z|$  so  $|z| - |w| \geq -|w - z|$ .

Like the absolute value in  $\mathbb{R}$ , we have by PM4  $|w - z| = |-1||z - w| = 1|z - w| = |z - w|$ , so if we combine the above two inequalities, we have  $||z| - |w|| \leq |z - w|$ .

Equivalently, using the same triangle from above, this follows from the fact that any one side of a triangle is longer than the difference of the other two sides.  $\square$

**C02.** Let  $a, b, c \in \mathbb{C}$ . Show that if  $\frac{b - a}{a - c} = \frac{a - c}{c - b}$  then  $|b - a| = |a - c| = |c - b|$ .

**C03.** Let  $n \geq 2$  be an integer. Prove that

$$\sum_{k=0}^{n-1} \cos\left(\frac{2k\pi}{n}\right) = 0 = \sum_{k=0}^{n-1} \sin\left(\frac{2k\pi}{n}\right)$$

*Proof* (with help from Ainsley, Kenson, Mabel). Let  $n \neq 1$  be a natural number. Then, we have that the  $n$ -th roots of unity are given by

$$\cos\left(\frac{2k\pi}{n}\right) + i \sin\left(\frac{2k\pi}{n}\right)$$

for  $k = 0, 1, 2, \dots, n - 1$ . Let  $z$  be the sum of the  $n$ -th roots of unity. Then,

$$z = \sum_{k=0}^{n-1} \left( \cos\left(\frac{2k\pi}{n}\right) + i \sin\left(\frac{2k\pi}{n}\right) \right)$$

The conclusion can equivalently be stated as that  $\operatorname{Re}(z) = 0$  and  $\operatorname{Im}(z) = 0$ . The only complex number that satisfies this is  $z = 0$ .

Now, let  $a = \cos\left(\frac{2\pi}{n}\right) + i \sin\left(\frac{2\pi}{n}\right)$ , the root of unity with  $k = 1$ . Then, we have that each root of unity is given by  $a^j$  for  $j = 1, 2, \dots, n$ . Since  $n \neq 1$ ,  $a = \operatorname{cis}\frac{2\pi}{n} \neq 1$  and  $z = 1 + a + a^2 + \dots + a^{n-1}$ .

Recall that the polynomial  $a^n - 1$  for  $n \geq 2$  factors as  $(a - 1)(a^{n-1} + a^{n-2} + \dots + a^2 + a + 1)$ . It follows that  $a^n - 1 = 1 - 1 = 0$  and  $0 = (a - 1)z$  so, from above,  $a \neq 1$  so  $z = 0$ .  $\square$